



The Impact of Preoperative Duration of Fasting on the Intravascular Volume Status of Children Older than 5 Years of Age: A Prospective, Observational Study

Beliz Bilgili, Tümay Umuroğlu

Marmara University Faculty of Medicine, Department of Anaesthesiology and Intensive Care, İstanbul, Türkiye

Cite this article as: Bilgili B, Umuroğlu T. The impact of preoperative duration of fasting on the intravascular volume status of children older than 5 years of age: a prospective, observational study. *Turk J Anaesthesiol Reanim.* 2025;53(5):226-232.

Abstract

Objective: Preoperative fasting is a common practice aiming to reduce the risk of pulmonary aspiration during anaesthesia. It is advised to avoid fasting times longer than 6 hours in all children, whenever possible. Prolonged fasting can be uncomfortable for children and may lead to dehydration and other negative outcomes. The primary outcome of the study was the relationship between preoperative duration of fasting and cardiac index (CI) variability, used as a surrogate for intravascular volume status after the induction of anaesthesia, in paediatric patients undergoing surgery.

Methods: Prospective, observational study that included patients over 5 years of age, scheduled for surgery. Passive leg-raising-induced CI variability was evaluated for fluid responsiveness and intravascular volume after anaesthesia induction. Patients were termed fluid responders (Rs) if an increase in CI of $>10\%$ was obtained after passive leg raising, and non-responders (NRs) if the CI variability was $<10\%$. CI and aortic peak velocity (V_{peak}) were measured through the suprasternal notch via an ultrasonic cardiac output monitor.

Results: There were 32 Rs and 53 (NRs). The mean duration of fasting for Rs was 11.53 ± 2.61 , while NR had a mean duration of fasting of 10.6 ± 2.93 hours, showing an insignificant difference. Aortic V_{peak} change was significantly higher in Rs (0.24 ± 0.17) compared to NRs (0.03 ± 0.13) ($P < 0.001$). Duration of fasting showed no significant correlation with CI variability and peak aortic velocity.

Conclusion: With this study method, it was observed that preoperative fasting time had no effect on intraoperative intravascular volume.

Keywords: Cardiac index, fasting time, fluid responsiveness, intravascular volume, paediatric anaesthesia, passive leg raising

Main Points

- Paediatric patients have higher insensible losses due to relatively increased surface area, increased respiratory rate, and greater metabolic rate, and are therefore more prone to become hypovolemic in the perioperative period.
- Prolonged fasting can be uncomfortable for children and may lead to dehydration and other negative outcomes.
- Even extensively prolonged preoperative fasting times had no significant impact on intravascular volume status in children over 5 years of age under anaesthesia.

Introduction

Preoperative fasting is a common practice aiming to reduce the risk of pulmonary aspiration during anaesthesia. Guidelines typically recommend specific fasting times including clear fluids up to 2 hours, breast milk up to 4 hours, and a light meal up to 6 hours before elective procedures.¹ It is advised to avoid prolonged fasting times longer than 6 hours in all children, whenever possible.² Nevertheless, the actual duration of fasting is frequently much longer than advised, including clear fluids abstinence.^{3,4} This undesirable situation is mostly inevitable and unpredictable, occurring because of delays in scheduling or overly cautious practices. Paediatric patients have higher insensible

losses due to increased surface area, increased respiratory rate, and greater metabolic rate, and are therefore more prone to become hypovolemic during the perioperative setting.⁵ Prolonged fasting can be uncomfortable for children and may lead to dehydration and other negative outcomes.⁶ Adequate hydration can prevent dehydration-related complications and delayed recovery.⁷

As prolonged fasting may lead to dehydration that may predispose paediatric patients to hemodynamic instability during anaesthesia it must be carefully managed. There may be a relationship between the duration of preoperative fasting and the occurrence of hypotension, which is a potential indicator of intravascular volume depletion. Fluid fasting times longer than 6 hours were associated with an increased risk of postinduction hypotension, during surgical preparation, although this association seemed to be non-linear.⁸ However, another study found no association between fasting duration and hypotension.⁹ The literature presents conflicting findings on this subject. Additionally, it should be noted that blood pressure is generally maintained in children with hypovolemia, and the lack of age-specific definitions for hypotension limits the ability of blood pressure to accurately reflect intravascular volume status.

The impact of preoperative fasting time on intravascular volume in paediatric patients remains a subject of debate and investigation. Understanding the impact of fasting time on intravascular volume status is crucial for optimizing perioperative care in paediatric patients. Recent research has focused on dynamic parameters and functional hemodynamic monitoring techniques to assess intravascular volume and fluid responsiveness. Since conventional static measures (e.g., blood pressure, urine output) may not accurately reflect real-time intravascular volume shifts, dynamic parameters such as cardiac index (CI) variability and aortic peak velocity (V_{peak}) serve as primary indicators of volume status. CI represents cardiac output (CO) normalized to body surface area, providing a better assessment of circulatory adequacy in paediatric patients than CO alone. Dynamic changes in CI following a preload challenge [such as passive leg raising (PLR)] offer insight into fluid responsiveness, which reflects intravascular volume status. Aortic V_{peak} serves as a non-invasive indicator of stroke volume variability.^{10,11} PLR-induced >10% CI variability reflects fluid responsiveness and intravascular volume deficit.¹¹ Changes in V_{peak} following PLR correlate with fluid responsiveness and preload dependency, making it an important parameter for assessing circulatory volume status.¹² PLR avoids unnecessary fluid administration by identifying true volume-responsive patients. It is particularly useful in paediatric patients, where baseline hemodynamic values may fluctuate, and traditional static markers are less predictive.¹³

This study tested the hypothesis that preoperative fasting time has an impact on intravascular volume. The primary outcome was the relation between preoperative fasting time and CI variability after anaesthesia, induction in paediatric patients.

Methods

Study Design

This prospective, observational, single-center trial was conducted after receiving approval from the Institutional Ethics Committee of Marmara University Faculty of Medicine Clinical Research (approval no.: 09.2017.669, date: 08.12.2017) in accordance with the principles outlined in the Declaration of Helsinki.

The trial was registered prior to patient enrollment at clinicaltrials.gov (<https://clinicaltrials.gov/>). Written informed consent was obtained from all patients' legal representatives included in the study. The flow chart of the study is shown in Figure 1.

Patients over 5 years of age and with American Society of Anesthesiologists (ASA) physical status I, scheduled as outpatients for surgery, were asked to enroll in the study. Paediatric patients, who were otherwise healthy except for the need for surgery and who did not receive intravenous (IV) fluid prior to anaesthesia induction, were included. Patients who failed to meet inclusion criteria and refused to give informed consent were excluded.

Anaesthesia Management

In the operating room, patients were monitored with an electrocardiogram, non-invasive blood pressure monitoring a pulse oximeter, monitors for anaesthetic gases, and a carbon dioxide (CO₂) analyzer. Anaesthesia, was initiated through an IV route, with 1 µg kg⁻¹ of remifentanyl hydrochloride, 5 mg kg⁻¹ of sodium thiopental, and 0.6 mg kg⁻¹ of rocuronium bromide. Following induction, the patient's airway was secured via tracheal intubation, and a maintenance dose of 2% sevoflurane in a mixture of 50% nitrous oxide and 50% oxygen was administered. Mechanical ventilation was controlled by the auto-flow mode of the Draeger Primus ventilator, with tidal volumes set at 8 mL per kg⁻¹ of body weight. The respiratory rate was adjusted to maintain end-tidal CO₂ between 35 and 45 mmHg, and a PEEP of 4 cm H₂O was used for respiratory support. To keep the patient warm throughout the surgery, the active air warming system was employed.

Protocol

After tracheal intubation, hemodynamic stability had been maintained with the aim of keeping a heart rate (HR) change of less than 10% for three minutes. Then, measurements were performed. The baseline measurements (T1) were recorded




Figure 1. Flow chart

with the patient in the semi-recumbent position. Next, the PLR maneuver was carried out by placing the patient in the supine position and simultaneously raising the patient's leg to 45°; after 1 minute (T2), a second set of measurements was recorded. In every stage, CI, CO, and aortic V_{peak} were measured; HR, CI, and mean arterial pressure were recorded. Cardiac measurement was performed through the suprasternal notch, via the Ultrasonic CO Monitor (USCOM) system by the same operator. Each parameter was measured consecutively three times, and the mean value of these three measurements was recorded.

An increase in CI (ΔCI) $>10\%$ following PLR is considered indicative of fluid responsiveness and suggests a functional intravascular volume deficit. Patients were termed fluid responders (Rs) if ΔCI of $>10\%$ was obtained after PLR, or non-responders (NRs) if ΔCI was $<10\%$.

Statistical Analysis

Data were evaluated for normality of distribution and reported as mean, standard deviation (SD), median, 25th and 75th interquartile, frequency, percentage, and minimum and maximum range. The independent t-test was employed to compare the means between two unrelated groups, and paired samples t-test was used to compare the means of two related groups. Qualitative data was compared using the Pearson chi-square test. Receiver operating characteristic (ROC) curves were built, sensitivity and specificity of variables were calculated for various values, and the value with the highest Youden index was taken as a cut-off point. Spearman correlation coefficient was used to test the association between variables. Statistical significance was defined as $P < 0.05$.

Sample size was calculated based on a previous adult study investigating the functional intravascular volume deficit in patients before surgery.¹⁴ Flow time corrected (FTc) data of

this study were taken into consideration. The FTc rise from $288 (\pm 45)$ ms to $316 (\pm 39)$ ms (before and after stroke volume maximization) was suspected to be significant. Accordingly, the minimum sample size required was 82 at an alpha level, and a beta level of 0.05 and 0.20, respectively. Based on possible dropouts, a total of 100 patients were planned for inclusion.

Results

As shown in Figure 1, a total of 100 paediatric patients with ASA I, scheduled for outpatient surgery, were initially assessed for eligibility. Fifteen patients were subsequently excluded due to either the preoperative administration of IV fluids or their declining participation. A total of 85 patients were included. There were 32 R and 53 NR. The patient characteristics of two groups were comparable in terms of age, gender, height, and body weight. The mean age of the participants was 7.82 ± 2.43 years. The mean fasting time was 10.95 ± 2.84 hours. The mean fasting time for Rs was 11.53 hours, with a SD of 2.61, while NR had a mean fasting time of 10.6 hours, with a SD of 2.93. This difference was not statistically significant, with a P value of 0.145 (Table 1).

The hemodynamic parameters before (T1) and after (T2) PLR are shown in Table 2. HR significantly decreased after PLR in both R and NR ($P < 0.001$, $P < 0.001$, respectively), but showed an insignificant difference between the study groups. Mean arterial pressure values were comparable among the groups at all study time points. CO was significantly lower in R than in NR in T1 ($P = 0.043$). CO significantly increased after PLR in R ($P < 0.001$) but remained similar in NR. ΔCO was significantly higher in R (0.89 ± 0.56) than in NR (-0.12 ± 0.49) ($P < 0.001$). V_{peak} was comparable among the groups in T1. Rs had a significantly increased V_{peak} in T2 ($P < 0.001$), and ΔV_{peak} was significantly

Table 1. Patient Characteristics

	Total (n = 85)	Responders (n = 32)	Non-responders (n = 53)	P value
	Mean \pm SD	Mean \pm SD	Mean \pm SD	
Age	7.82 \pm 2.43	7.69 \pm 2.61	7.91 \pm 2.33	^a 0.690
Gender n (%)				^b 0.637
Female	27 (100)	9 (33.3)	18 (66.7)	
Male	58 (100)	23 (39.7)	35 (60.3)	
Height	124.36 \pm 18.11	124.59 \pm 20.54	124.23 \pm 16.68	^a 0.928
Weight	27.33 \pm 9.72	29.59 \pm 11.84	25.96 \pm 7.99	^a 0.131
Fasting time	10.95 \pm 2.84	11.53 \pm 2.61	10.6 \pm 2.93	^a 0.145

a, Independent t-test; b, pearson chi-square test. SD, standard deviation

Table 2. Hemodynamic Parameters Recorded Before (T1) and After (T2) PLR

Mean \pm SD		Responders (n = 32)	Non-responders (n = 53)	^a P (inter group)
		Mean \pm SD		
HR	T1	110.63 \pm 20.05	110.04 \pm 14.66	0.877
	T2	98.38 \pm 17.03	96.15 \pm 14.27	0.519
	Δ	-12.25 \pm 8.93	-13.89 \pm 7.83	0.378
	^c P (intra group)	< 0.001**	< 0.001**	
MAP	T1	71.38 \pm 12.6	69.26 \pm 8.27	0.403
	T2	69.16 \pm 13.21	68.98 \pm 7.18	0.945
	Δ	-2.22 \pm 9.48	-0.28 \pm 6.63	0.272
	^c P	0.195	0.757	
CI	T1	2.91 \pm 0.81	3.58 \pm 0.94	0.001**
	T2	3.8 \pm 1.08	3.44 \pm 0.82	0.092
	Δ	0.89 \pm 0.53	-0.13 \pm 0.44	< 0.001**
	^c P	< 0.001**	0.032*	
CO	T1	2.93 \pm 0.81	3.39 \pm 1.12	0.043*
	T2	3.81 \pm 1.16	3.27 \pm 0.97	0.023*
	Δ	0.89 \pm 0.56	-0.12 \pm 0.49	< 0.001**
	^c P	< 0.001**	0.076	
V _{peak}	T1	0.93 \pm 0.25	1.02 \pm 0.24	0.101
	T2	1.17 \pm 0.27	1.05 \pm 0.21	0.029*
	Δ	0.24 \pm 0.17	0.03 \pm 0.13	< 0.001**
	^c P	< 0.001**	0.055	

a, Independent t-test; c, Paired Samples t-test, *P < 0.05, **P < 0.01

PLR, passive leg raising; HR, heart rate; MAP, mean arterial pressure; CI, cardiac index; CO, cardiac output; V_{peak}, aortic peak velocity Δ : change; SD, standard deviation

higher in Rs (0.24 \pm 0.17) compared to NRs (0.03 \pm 0.13) (P < 0.001). The optimal cut-off value for ΔV_{peak} is >0.17, which corresponds to the highest Youden index of 0.63. It has a sensitivity of 84.9% and specificity of 78.1%. The area under ROC (AuROC) curve is 0.872 [95% CI (0.784, 0.961), P <

0.001]. Upon evaluating the relationships between fasting time and various cardiac measurements, including CI, CO, and V_{peak}, fasting time showed no significant correlation with parameters (Table 3).

Table 3. Correlation Between Fasting Time and Cardiac Measurements

	Δ CI	Δ CO	ΔV_{peak}
Fasting time			
r	0.117	0.126	0.199
*P	0.285	0.249	0.067
a, Independent t-test; r, pearson correlation coefficient; Δ CI, change in cardiac index; Δ CO, change in cardiac output; ΔV_{peak} , change in aortic peak velocity			

Discussion

The objective of this study was to evaluate the impact of preoperative fasting duration on intravascular volume status in paediatric patients undergoing elective surgery. Using dynamic hemodynamic assessment via PLR, we found that prolonged fasting had no significant effect on intravascular volume status, as determined by CI variability (Δ CI), CO, and aortic V_{peak} .

The actual duration of preoperative fasting in paediatric patients is often longer than current recommendations, varying between 11-13 hours.¹⁵⁻¹⁷ Guidelines suggest reducing the clear fluid fasting period to approximately 2 hours before surgery to minimize discomfort and lower the incidence of hypotension in the perioperative period. However in clinical practices, paediatric fasting guidelines are often not applicable, resulting in longer fasting periods than recommended, with a high incidence of discomfort due to hunger and thirst.¹⁶ The present study, which found an average fasting time of 11 hours fasting time revealed similar prolonged fluid abstinence times in children scheduled for outpatient surgery.

In this study, the patients having more than 10% increase in CI following PLR had comparable fasting times with NRs. There was no correlation between the duration of fasting and CO and V_{peak} . A study has suggested that prolonged fasting (more than 6 hours for liquid and 12 hours for solid) may increase the risk of hypotension during anaesthesia induction, particularly in paediatric patients.⁹ A retrospective analysis was conducted focusing on clear fluids fasting times in children after implementing a 1 hour clear fasting rule. This study noted that longer fasting times, particularly those significantly exceeding the recommendations of guidelines, were associated with adverse events related to hypovolemia dehydration, such as difficulty in vein cannulations, electrocardiogram alterations, and episodes of hypotension.¹⁶ Our findings contrast with prior studies that have reported associations between fasting duration and intraoperative hypotension. However, these studies primarily evaluated blood pressure changes as a surrogate for intravascular volume depletion, without incorporating dynamic assessments of fluid responsiveness.

Dynamic parameters, rather than static ones, have proven to be more reliable for predicting volume responsiveness in mechanically ventilated children.¹⁰ In contrast to static indicators of preload, dynamic tests analyze the effect of a defined change in preload on stroke volume or CO, thus giving an indication of the patient's position on the Frank-Starling curve. Dynamic indices, therefore, are better predictors of changes in preload on the steep portion of the Frank-Starling curve and have been validated in predicting the response to volume expansion in both adults and children.^{18,19} CI changes following PLR can identify patients who will benefit from fluid administration.²⁰ The effectiveness of the PLR test partly depends on the ability to significantly alter venous return by elevating the legs. In smaller children, particularly in infants and toddlers, the relative volume shift achieved by lifting the legs may not significantly change the preload due to their smaller blood volume and different body proportions. PLR induced CI changes reliably reflects fluid responsiveness in children older than 5 years of age,²⁰ but under this group of age it cannot identify all Rs with certainty.¹³ This is the reason why study included children older than 5 years of age. Also, aortic ΔV_{peak} predicted fluid responsiveness with an AuROC of 0.872 in the present study, supporting the evidence that aortic blood flow velocity is a reliable index to predict fluid responsiveness in children.¹¹ USCOM measuring CO via a probe applied to the suprasternal notch revealed stroke volume variation as a reliable index in the prediction of fluid responsiveness.²¹ The change in V_{peak} measured by USCOM showed a significant correlation with the measurements obtained through the apical 5-chamber view.²² In light of these findings, we measured dynamic indices using USCOM.

Given the similar fasting times of Rs and NRs, it was suggested that fluid responsiveness cannot be directly related to the abstinence of oral intake. One-third of the children in our study were resistant; the reason for this could be attributed to the vasodilatory effects of the anaesthetic agents during the induction period. CO is directly related to the so-called stressed volume of the circulation.²³ For optimal cardiac preload, it's essential to have an adequate portion of the blood volume under stress, which depends on the state of the intravascular volume (normovolemia) and the tension within the blood vessels (vasotension).²⁴ One potential explanation for the lack of correlation between fasting duration and hemodynamic changes is anaesthesia-induced vasodilation, which significantly alters intravascular dynamics independent of fasting status. General anaesthesia induction and neuraxial anaesthesia lead to vasodilation, reducing the stressed volume and causing a functional intravascular deficit.^{14,25} The presence of a functional intravascular deficit could lead to changes in the CI when a PLR test is performed. Muller et al.²⁷ showed that blood volume assessed by a dynamic test did not decrease after

preoperative fasting. As measurements were performed before anaesthesia, hemodynamics were not affected by anaesthesia induction. At this point, our study assesses the impacts on both the duration of fasting and anaesthesia by evaluating dynamic parameters following anaesthesia induction. Thus, in our study population, any fasting-related reduction in circulating volume may have been overshadowed by the hemodynamic effects of anaesthesia, explaining the absence of a significant correlation between fasting time and fluid responsiveness.

Children exhibit unique physiological adaptations that allow them to maintain hemodynamic stability despite fasting-related changes in intravascular volume. Unlike adults, where stroke volume plays a dominant role in CO regulation, young children primarily rely on HR adjustments to preserve CO. In paediatric patients, stroke volume is relatively fixed due to less compliant ventricles, meaning HR plays the primary role in adjusting CO²³. Even if fasting led to mild intravascular depletion, children may compensate by increasing HR, thereby preserving CO and preventing a detectable change in CI or CO.¹⁹

Study Limitations

While our study provides important insights into fasting and hemodynamics in paediatric patients, certain limitations should be acknowledged. Primarily, it was conducted in a single institution, which may affect the generalizability of the findings. Additionally, the study included only children who were older than 5 years and were otherwise healthy apart from their surgical requirements. This selection criterion may restrict the applicability of our results to younger children, such as infants and toddlers, as well as to children with other comorbidities. Our study assessed volume status after anaesthesia induction, which means we could not evaluate fasting-induced changes before anaesthesia. While we assessed intravascular volume status, we did not measure interstitial fluid shifts, which could also play a role in perioperative hemodynamics.

Future studies may focus on a wider age range of children and those with cardiac, renal, and pulmonary comorbidities to assess the impact of fasting time on intravascular volume status more comprehensively and monitor longitudinal hemodynamic parameters before and after anaesthesia induction.

Conclusion

In conclusion, preoperative fasting time had no effect on intraoperative intravascular volume. Anaesthesia-induced vasodilation and paediatric compensatory mechanisms (HR-driven CO maintenance, sympathetic activation) likely mitigate any fasting-related hemodynamic effects, explaining

the lack of correlation between fasting time and fluid responsiveness. These findings emphasize the importance of using dynamic assessments over static indicators when evaluating perioperative fluid status and highlight the need for further research on fasting and hemodynamic adaptations in paediatric patients.

Ethics

Ethics Committee Approval: The ethical approval was obtained from the Institutional Ethics Committee of Marmara University Faculty of Medicine Clinical Research (approval no.: 09.2017.669, date: 08.12.2017).

Informed Consent: Written informed consent was obtained from all patients' legal representatives included in the study.

Footnotes

Author Contributions: Surgical and Medical Practices - B.B., T.U.; Concept - B.B., T.U.; Design - B.B., T.U.; Data Collection and/or/ Processing - B.B., T.U.; Analysis and/or/Interpretation - B.B., T.U.; Literature Review - B.B., T.U.; Writing - B.B., T.U.

Declaration of Interests: The authors declare no conflicts of interest.

Funding: No funding was received for conducting this study.

References

1. Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: Application to healthy patients undergoing elective procedures: An updated report by the American Society of Anesthesiologists Task Force on preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration. *Anesthesiology*. 2017;126(3):376-393. [\[CrossRef\]](#)
2. Frykholm P, Disma N, Andersson H, et al. Pre-operative fasting in children: a guideline from the European Society of Anaesthesiology and Intensive Care. *Eur J Anaesthesiol*. 2022;39(1):4-25. [\[CrossRef\]](#)
3. Joshi GP, Abdelmalak BB, Weigel WA, et al. 2023 American Society of Anesthesiologists Practice Guidelines for preoperative fasting: carbohydrate-containing clear liquids with or without protein, chewing gum, and pediatric fasting duration-a modular update of the 2017 American Society of Anesthesiologists Practice Guidelines for preoperative fasting. *Anesthesiology*. 2023;138(2):132-151. [\[CrossRef\]](#)
4. Dolgun E, Yavuz M, Eroğlu B, Islamoğlu A. Investigation of preoperative fasting times in children. *J Perianesth Nurs*. 2017;32(2):121-124. [\[CrossRef\]](#)
5. Kight BP, Waseem M. Pediatric fluid management. *StatPearls*. Treasure Island (FL): StatPearls Publishing
6. Copyright © 2024, StatPearls Publishing LLC; 2024. [\[CrossRef\]](#)
7. Buller Y, Sims C. Prolonged fasting of children before anaesthesia is common in private practice. *Anaesth Intensive Care*. 2016;44(1):107-110. [\[CrossRef\]](#)
8. Derieg S. An overview of perioperative care for pediatric patients. *Aorn J*. 2016;104(1):4-10. [\[CrossRef\]](#)
9. Simpao AF, Wu L, Nelson O, et al. Preoperative fluid fasting times and postinduction low blood pressure in children: a retrospective analysis. *Anesthesiology*. 2020;133(3):523-533. [\[CrossRef\]](#)
10. Hajian P, Shabani M, Khanlarzadeh E, Nikooseresht M. The impact of preoperative fasting duration on blood glucose and hemodynamics in children. *J Diabetes Res*. 2020;2020:6725152. [\[CrossRef\]](#)

11. Yenjabog P, Kanchongkitiphon W, Chutipongtanate S, Lertbunrian R, Ungprasert P. Dynamic parameters for fluid responsiveness in mechanically ventilated children: a systematic review. *Front Pediatr*. 2022;10:1010600. [\[CrossRef\]](#)
12. Gan H, Cannesson M, Chandler JR, Ansermino JM. Predicting fluid responsiveness in children: a systematic review. *Anesth Analg*. 2013;117(6):1380-1392. [\[CrossRef\]](#)
13. Sethasathien S, Jariyasakoolroj T, Silvilairat S, Srisurapanont M. Aortic peak flow velocity as a predictor of fluid responsiveness in mechanically ventilated children: a systematic review and meta-analysis. *Pediatr Crit Care Med*. 2023;24(7):e352-e361. [\[CrossRef\]](#)
14. El-Nawawy AA, Farghaly PM, Hassouna HM. Accuracy of passive leg raising test in prediction of fluid responsiveness in children. *Indian J Crit Care Med*. 2020;24(5):344-349. [\[CrossRef\]](#)
15. Bundgaard-Nielsen M, Jørgensen CC, Secher NH, Kehlet H. Functional intravascular volume deficit in patients before surgery. *Acta Anaesthesiol Scand*. 2010;54(4):464-469. [\[CrossRef\]](#)
16. Tsukamoto M, Hitosugi T, Yokoyama T. Influence of fasting duration on body fluid and hemodynamics. *Anesth Prog*. 2017;64(4):226-229. [\[CrossRef\]](#)
17. Aroonpruksakul N, Punchuklang W, Kasikan K, et al. The actual duration of preoperative fasting in pediatric patients, and its effects on hunger and thirst: a prospective observational study. *Transl Pediatr*. 2023;12(2):146-154. [\[CrossRef\]](#)
18. Li Y, Lu Q, Wang B, Tang W, Fan L, Li D. Preoperative fasting times for patients undergoing elective surgery at a pediatric hospital in Shanghai: the big evidence-practice gap. *J Perianesth Nurs*. 2021;36(5):559-563. [\[CrossRef\]](#)
19. Monnet X, Marik PE, Teboul JL. Prediction of fluid responsiveness: an update. *Ann Intensive Care*. 2016;6(1):111. [\[CrossRef\]](#)
20. Lee JH, Kim EH, Jang YE, Kim HS, Kim JT. Fluid responsiveness in the pediatric population. *Korean J Anesthesiol*. 2019;72(5):429-440. [\[CrossRef\]](#)
21. Lukito V, Djer MM, Pudjiadi AH, Munasir Z. The role of passive leg raising to predict fluid responsiveness in pediatric intensive care unit patients. *Pediatr Crit Care Med*. 2012;13(3):e155-160. [\[CrossRef\]](#)
22. Cheng YW, Xu F, Li J. Identification of volume parameters monitored with a noninvasive ultrasonic cardiac output monitor for predicting fluid responsiveness in children after congenital heart disease surgery. *Medicine (Baltimore)*. 2018;97(39):e12289. [\[CrossRef\]](#)
23. Devauchelle P, de Queiroz Siqueira M, Lilot M, Chassard D, Desgranges FP. Suprasternal notch echocardiography: a potential alternative for the measurement of respiratory variation in aortic blood flow peak velocity in mechanically ventilated children. *J Clin Monit Comput*. 2018;32(3):589-591. [\[CrossRef\]](#)
24. Jacob M, Chappell D. Effects of perioperative fasting on haemodynamics and intravascular volumes. *Best Pract Res Clin Anaesthesiol*. 2012;26(4):421-430. [\[CrossRef\]](#)
25. Maas JJ, Pinsky MR, Aarts LP, Jansen JR. Bedside assessment of total systemic vascular compliance, stressed volume, and cardiac function curves in intensive care unit patients. *Anesth Analg*. 2012;115(4):880-887. [\[CrossRef\]](#)
26. Bundgaard-Nielsen M, Holte K, Secher NH, Kehlet H. Monitoring of peri-operative fluid administration by individualized goal-directed therapy. *Acta Anaesthesiol Scand*. 2007;51(3):331-340. [\[CrossRef\]](#)
27. Muller L, Brière M, Bastide S, et al. Preoperative fasting does not affect haemodynamic status: a prospective, non-inferiority, echocardiography study. *Br J Anaesth*. 2014;112(5):835-841. [\[CrossRef\]](#)